

Please write clearly in block capitals.

Centre number

Candidate number

Surname

Forename(s)

Candidate signature

I declare this is my own work.

# INTERNATIONAL A-LEVEL PHYSICS

## Unit 5 Physics in practice

Thursday 18 January 2024

07:00 GMT

Time allowed: 2 hours

### Materials

For this paper you must have:

- a Data and Formulae Booklet as a loose insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate
- a protractor.

### Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each question or on blank pages.
- All working must be shown.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

### Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
<b>TOTAL</b>	



**Section A**Answer **all** questions in this section.**0 1 . 1**

Describe how to reduce the effect of random errors on a set of measurements.

**[1 mark]**


---



---



---

A student measures the length  $l$  of a simple pendulum six times using a metre ruler.**Table 1** shows her measurements.**Table 1**

$l / \text{cm}$	89.4	89.3	89.5	89.4	89.4	89.5
-----------------	------	------	------	------	------	------

**0 1 . 2**Calculate the mean value of  $l$ .**[1 mark]**mean value of  $l =$  \_\_\_\_\_ cm

The student makes repeated measurements of the time for ten oscillations of the simple pendulum.

She calculates that the mean period for one oscillation is  $1.95 \text{ s} \pm 1\%$ .**0 1 . 3**

Comment on whether the student's measurements of time are precise.

**[1 mark]**


---



---



---



The student uses her measurements to determine a value for  $g$ .

0 1 . 4

Deduce whether the student's measurements are accurate.

[2 marks]

---

---

---

0 1 . 5

The percentage uncertainty in  $l$  is 0.1%.

Another student suggests that there is a systematic error in some of the data.

Discuss the evidence for this suggestion.

[1 mark]

---

---

---

6


Turn over for the next question

Turn over ►



0 2

Two students **A** and **B** used the apparatus shown in **Figure 1** to determine the count rate from radioactive sources. The activities of the sources did not vary during the determination.

**Figure 1**

Both students used the same apparatus and placed the detector 10 cm from the source. Each student measured the radioactive count for a time  $t$ .

**Table 2** shows the measurements made by the two students.

**Table 2**

	Count	$t / \text{s}$
student <b>A</b>	464	$60 \pm 1$
student <b>B</b>	2150	$300 \pm 1$

When a count is measured as  $N$ , the absolute uncertainty in the measurement is  $\sqrt{N}$ .

Each student calculated the count rate  $C$  using the equation:

$$C = \frac{N}{t}$$

The uncertainty in the measurement of the separation of the source and the detector is negligible.



0	2	.	1
---	---	---	---

State **two** reasons why student **A**'s determination of  $C$  has a greater percentage uncertainty than student **B**'s determination of  $C$ .

[2 marks]

1 \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

2 \_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

0	2	.	2
---	---	---	---

Deduce, by considering uncertainties, whether the data suggest that both students used the same radioactive source in their determinations of  $C$ .

[5 marks]

\_\_\_\_\_

\_\_\_\_\_

7
---

Turn over ►



**Figure 2** shows apparatus used to determine the number of lines per millimetre on a plane transmission diffraction grating. The laser emits light of wavelength 633 nm.

Diagram illustrating the setup for observing diffraction patterns:

- A **laser** beam is directed towards a **diffraction grating**.
- The **diffraction grating** is positioned between the laser and the **screen**.
- The **screen** is located to the right of the diffraction grating, where the diffraction pattern is observed.

In your answer you should:

- [6 marks]**

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and extend across the width of the page. There are no margins, text, or other markings on the paper.

*Do not write  
outside the  
box*

6


**Turn over for the next question**

**Turn over ►**



0 4

**Figure 3** shows apparatus used to investigate how the current  $I$  in a battery varies with the length  $l$  of resistance wire connected across its terminals. The battery has an emf  $\varepsilon$  of 3.0 V and an internal resistance  $r$ .

**Figure 3**

**Table 3** shows values of  $l$ ,  $I$  and  $I^{-1}$ .

**Table 3**


$l / \text{m}$	$I / \text{A}$	$I^{-1} / \text{A}^{-1}$
0.100	2.38	0.42
0.300	1.02	0.98
0.500	0.72	1.39
0.700	0.52	1.92
0.900	0.44	2.27

0 4 . 1

Draw, on **Figure 4**, a graph to show the variation of  $I^{-1}$  with  $l$ . Draw a best-fit line.

**[4 marks]**



**Figure 4****Question 4 continues on the next page****Turn over ►**

**0 4 . 2**Determine the gradient of the graph that you have drawn in **Figure 4**.**[2 marks]**gradient = \_\_\_\_\_  $\text{A}^{-1} \text{m}^{-1}$  $R$  is the resistance of a length  $l$  of the wire.The equation that relates  $I$  to  $R$  is:

$$\frac{1}{I} = \frac{R}{\varepsilon} + \frac{r}{\varepsilon}$$

**0 4 . 3**

The resistance wire has a diameter of 0.457 mm.

Calculate, using your answer to Question **04.2**, the resistivity of the material in the resistance wire.**[3 marks]**resistivity = \_\_\_\_\_  $\Omega \text{m}$ **0 4 . 4**Calculate  $r$  using data from your graph in **Figure 4**.**[2 marks]** $r =$  \_\_\_\_\_  $\Omega$ **11**

**Turn over for the next question**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**




0 5

Astronomers have measured the orbital period  $T$  and the orbital radius  $r$  for five planets in the solar system of the star HD 108236.

**Figure 5** shows the variation of  $\ln(T / \text{s})$  with  $\ln(r / \text{m})$  for the astronomers' data.

**Figure 5**



Theory suggests that the relationship between  $T$  and  $r$  is:

$$T^2 = \frac{4\pi^2 r^3}{GM}$$

where  $M$  is the mass of the star HD 108236.



- 0 5 . 1** Show that the intercept  $c$  on a graph of  $\ln(T / \text{s})$  against  $\ln(r / \text{m})$  is given by:

$$c = \frac{1}{2} \ln \left( \frac{4\pi^2}{GM} \right)$$

**[2 marks]**

- 0 5 . 2** Explain, without calculation, how the gradient of the graph in **Figure 5** can be used to show that the astronomers' data are consistent with the relationship on page 12.

**[1 mark]**

---



---



---

- 0 5 . 3** Determine  $M$ .

**[3 marks]**

$M =$  \_\_\_\_\_ kg

**END OF SECTION A**


**Turn over ►**



## Section B

Answer **all** questions in this section.**0 6**

**Figure 6** shows the apparatus used by Joule to measure the effect of doing work and to determine the specific heat capacity of water.

**Figure 6**

**P** and **Q** are masses each of mass  $M$ . They fall from rest from an initial height  $h$  above the ground. They are connected to a pulley system that turns sets of paddles. The paddles do work by stirring the water in the container.

The gravitational potential energies of **P** and **Q** are transferred to the water. The temperature of the water increases as a result of the work done as **P** and **Q** fall.

When **P** and **Q** reach the ground, the apparatus is reset as in **Figure 6** without causing the paddles to turn. **P** and **Q** are allowed to fall a total of 20 times.

The data collected are:

$$\begin{aligned} h &= 1.5963 \text{ m} \\ M &= 13.158 \text{ kg} \\ m &= 6.2512 \text{ kg} \\ \Delta\theta &= 0.364 \text{ K} \end{aligned}$$

where  $\Delta\theta$  is the total increase in temperature of the water and  $m$  is the mass of the water.



**06.1**

The initial temperature of the water in the container is lower than the temperature of the surroundings.

Explain, with reference to the first law of thermodynamics, why the temperature of the water increases during this experiment.

**[3 marks]**

---

---

---

---

---

---

---

---

**06.2**

Show that the total energy transferred by **P** and **Q** as they fall is approximately 8240 J.

**[2 marks]****06.3**

State and explain the appropriate number of significant figures that should be given in the answer to Question **06.2**.

**[1 mark]**

---

---

**Question 6 continues on the next page**

**Turn over ►**

After each release, **P** and **Q** accelerate briefly before falling at a constant speed of  $6.15 \text{ cm s}^{-1}$  until they hit the ground.

0 6 . 4

Explain, in terms of the forces acting on the paddles, why **P** and **Q** accelerate briefly before falling at constant speed.

[2 marks]

---

---

---

---

---

---

0 6 . 5

Calculate the total kinetic energy of **P** and **Q** before they hit the ground.

[1 mark]

total kinetic energy = \_\_\_\_\_ J

0 6 . 6

Explain whether the work done on the water by **P** and **Q** is significantly affected by the total kinetic energy of **P** and **Q**.

[2 marks]

---

---

---





0	6	.	7
---	---	---	---

The total energy transferred by **P** and **Q** as they fall is approximately 8240 J.

$$\Delta\theta = 0.364 \text{ K}$$

$$m = 6.2512 \text{ kg}$$

Joule estimated that:

- 360 J was used to increase the temperature of the container and the paddles
- 0.060 K of the total increase in temperature of the water was caused by thermal conduction from the surroundings.

Calculate a value for the specific heat capacity of water.

**[3 marks]**

specific heat capacity = \_\_\_\_\_ J kg<sup>-1</sup> K<sup>-1</sup>

**Question 6 continues on the next page**


**Turn over ►**



**Figure 7** shows the diameters of the pulleys and the drum. The pulleys are frictionless. The container and the water have been omitted from **Figure 7**.


**P** and **Q** each has a mass  $M$  of 13.158 kg and produce tensions in their strings of  $T_P$  and  $T_Q$  respectively.

**Figure 7**



**Figure 8** shows the strings and the drum viewed from above. These strings have tensions of  $T_1$  and  $T_2$  that together exert a couple on the drum.

**Figure 8**



**0 6 . 8** **P** and **Q** fall at a constant speed.

Show that the torque applied to the drum by the couple is approximately 0.9 N m.

**[3 marks]**



0	6	.	9
---	---	---	---

**P** and **Q** fall through height  $h$  a total of 20 times in 519 s.

The total work done on the drum by  $T_1$  and  $T_2$  during this time is 8240 J.

Calculate the angular velocity of the drum.

**[2 marks]**

angular velocity = \_\_\_\_\_  $\text{rad s}^{-1}$

19
----


**Turn over for the next question**

**Turn over ►**



0 7

**Figure 9** shows a capacitor **C** with wires attached to its two circular conducting plates **P** and **Q**. The plates are separated by an air gap of width  $d$ .

**Figure 9**

**P** and **Q** each has a radius of 1.3 cm. The capacitance of **C** is 52 pF.

0 7 . 1


Show that  $d$  is approximately  $9.0 \times 10^{-5}$  m.

**[2 marks]**

**Figure 10** shows **C** connected in a circuit with a resistor **R**, a switch **S** and a battery. The battery has an emf of 100 V and negligible internal resistance.

**R** has a resistance of  $820\text{ G}\Omega$ .

**Figure 10**



0 7 . 2

**C** is initially uncharged.  
**S** is then closed.

After a time  $t$ , the voltage  $V$  across **C** is 99.2 V.

Calculate  $t$ .

**[2 marks]**

$t =$  \_\_\_\_\_ s

**Question 7 continues on the next page**

**Turn over ►**



**S** remains closed and **C** becomes fully charged.

**C** is now used as a microphone.

A sound wave of frequency 100 Hz arrives at plate **P**. This causes **P** to oscillate while **Q** does not move.

**P** oscillates with simple harmonic motion at a frequency of 100 Hz and with an amplitude of  $4.5 \times 10^{-6}$  m.

The width of the air gap repeatedly increases and decreases.

**07.3**

Explain why the charge on **C** stays approximately constant as **P** oscillates.

[2 marks]

---

---

---

---

---

---

---

**07.4**

The width of the air gap decreases by  $4.5 \times 10^{-6}$  m compared with its value in Question **07.1**.

Deduce the effect of this change on  $V$ .

[3 marks]

---

---

---

---



0 7 . 5


The sound wave arrives at **P** at time  $t = 0$

At  $t = 0$ ,  $V = 100 \text{ V}$  and **P** begins to move towards **Q**.

Draw, on **Figure 11**, a graph to show the variation of  $V$  with  $t$  for **one** oscillation of **P**.  
Add axes and scales to your graph.

[3 marks]

Figure 11



12


Turn over for the next question

Turn over ►



0 8

**Figure 12** shows how the intensity of the radiation observed on Earth from a star varies with the frequency  $f$  of the radiation.

**Figure 12**

$\lambda_{\text{peak}}$  is the wavelength, in m, of the peak intensity of the radiation emitted from the surface of a star.

$\lambda_{\text{peak}}$  is related to the surface temperature  $T$ , in K, of the star by:

$$\lambda_{\text{peak}} = \frac{2.90 \times 10^{-3}}{T}$$

0 8 . 1

Calculate  $T$ .

**[2 marks]**

$T =$  \_\_\_\_\_ K





**Question 8 continues on the next page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Turn over ►**



0 8 . 2

White light is emitted from the interior of the star. Atoms in the star's outer layers absorb photons of certain frequencies. This causes the dips in intensity in **Figure 12**. The atoms in the outer layers are excited when they absorb the photons.

Explain why the photon frequencies at which the dips occur depend on the particular elements present in the outer layers of the star.

[2 marks]

---



---



---



---



---



---

Antares is a star system that is 550 light years from the Earth. A light year is the distance travelled by light in one year.

Antares emits  $7.59 \times 10^4$  times as much energy per second as the Sun.

0 8 . 3

Show that one light year is approximately equivalent to  $9.5 \times 10^{15}$  m.

[1 mark]

0 8 . 4

$I_A$  is the intensity of light from Antares at the position of the Earth.

$I_S$  is the intensity of light from the Sun at the position of the Earth.

Calculate  $\frac{I_A}{I_S}$

radius of the Earth's orbit around the Sun =  $1.5 \times 10^{11}$  m


[3 marks]

$$\frac{I_A}{I_S} = \underline{\hspace{5cm}}$$



**Figure 13** shows a binary system that consists of two stars Antares **A** and Antares **B**. The stars have circular orbits around their common centre of mass **C**. The gravitational forces  $F$  between the two stars and their distances from **C** are shown.

**Figure 13**



mass of **A** =  $2.39 \times 10^{31}$  kg

mass of **B** =  $9.95 \times 10^{30}$  kg

0 8 . 5

Calculate the time taken for Antares **A** to complete one orbit around **C**.

[4 marks]

time = \_\_\_\_\_ s

0 8 . 6

Explain why Antares **A** and Antares **B** must have the same orbital periods around **C**.

[1 mark]

---



---



---

END OF QUESTIONS



**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**



*Do not write  
outside the  
box*

[illegible]

*Do not write  
outside the  
box*

[illegible]

[illegible]

**There are no questions printed on this page**

*Do not write  
outside the  
box*

**DO NOT WRITE ON THIS PAGE  
ANSWER IN THE SPACES PROVIDED**

**Copyright information**

For confidentiality purposes, all acknowledgements of third-party copyright material are published in a separate booklet. This booklet is published after each live examination series and is available for free download from [www.oxfordaqa.com](http://www.oxfordaqa.com)

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright-holders may have been unsuccessful and OxfordAQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team.

Copyright © 2024 OxfordAQA International Examinations and its licensors. All rights reserved.

