

Please write clearly in block capitals.

Centre number

--	--	--	--	--

Candidate number

--	--	--	--

Surname _____

Forename(s) _____

Candidate signature _____

I declare this is my own work.

INTERNATIONAL AS PHYSICS

Unit 2 Electricity, waves and particles

Tuesday 21 May 2024

07:00 GMT

Time allowed: 2 hours

Materials

For this paper you must have:

- a Data and Formulae Booklet as a loose insert
- a ruler with millimetre measurements
- a scientific calculator, which you are expected to use where appropriate
- a protractor.

Instructions

- Use black ink or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- You must answer the questions in the spaces provided. Do not write outside the box around each question or on blank pages.
- All working must be shown.
- If you need extra space for your answer(s), use the lined pages at the end of this book. Write the question number against your answer(s).
- Do all rough work in this book. Cross through any work you do not want to be marked.

Information

- The marks for questions are shown in brackets.
- The maximum mark for this paper is 80.

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12–25	
TOTAL	



Section AAnswer **all** questions in this section.**0 1**

A simple pendulum completes 39 oscillations in 1.0 minute.

Calculate the length of the pendulum.

[2 marks]

length = _____ m

2**0 2**

Monochromatic light is incident on a clean metal surface and photoelectrons are emitted from the surface.

A photon of this light has energy of 4.5×10^{-19} J.

The stopping potential of the electrons is 1.1 V.

Calculate, in eV, the work function of the metal.


[2 marks]

work function = _____ eV

2

0 3

Figure 1 shows a cell in a series circuit with a load resistor. The cell has an internal resistance r .

Figure 1

The emf ε of the cell is given by

$$\varepsilon = IR + Ir$$

where I is the current in the circuit and R is the load resistance.

Explain how this equation is consistent with the law of conservation of energy.

[2 marks]

2

Turn over for the next question

Turn over ►




0 4

Figure 2 shows the I – V characteristic for a component.

Do not write
outside the
box

Figure 2



Calculate the resistance of the component when the current in it is 0.80 mA.

[2 marks]

resistance = _____ Ω

2



0 5

An electrical toy normally needs six cells.

Figure 3a shows the inside of the cell compartment of the toy. The markings show how the cells **A** to **F** should be arranged.

Figure 3b shows the circuit diagram of the cells when they are placed correctly into the compartment.

Figure 3a





Figure 3b




0 5 . 1

Draw lines on **Figure 4** to show how the cells should be connected in the compartment.

[1 mark]

Figure 4



Question 5 continues on the next page

Turn over ►



Each cell has an emf of 1.50 V and an internal resistance of $0.25\ \Omega$.

0 5 . 2

On one occasion, a child fits five cells into the toy but leaves out cell **B**.

Suggest and explain the consequences of leaving out cell **B**.


[2 marks]



0 5 . 3

On another occasion, a fault occurs that causes a connection between **X** and **Y** as shown in **Figure 5**. All six cells are fitted correctly. The connection has zero resistance.

Figure 5



Calculate the current in the connection between **X** and **Y**.

[3 marks]

current = _____ A

6

Turn over for the next question

Turn over ►



0	6
---	---

A sodium lamp contains low-pressure sodium gas.
An electric current in the sodium gas causes the sodium atoms to emit light.

0	6	.	1
---	---	---	---


Explain how the sodium atoms become excited.

[2 marks]



Figure 6 shows part of the line spectrum for the light emitted from the sodium lamp.

Figure 6



0 6 . 2

Explain why the spectrum produced by the sodium lamp has light of only certain wavelengths.

[3 marks]

0 6 . 3

Calculate, in J, the energy of the **least** energetic photon that corresponds to a spectral line in **Figure 6**.

[3 marks]

photon energy = _____ J

8

Turn over ►




0 7

Figure 7 shows apparatus used to investigate stationary waves in a column of air inside a cylindrical tube.

A loudspeaker, connected to a frequency generator, emits a sound wave with a single frequency into the open end of the tube.

Figure 7



The sound wave from the loudspeaker travels through the air in the tube and reflects at the surface of the water.

0 7 . 1

The frequency of the sound is increased from zero. At a certain frequency f , the loudness of the sound in the tube increases significantly.

Explain, with reference to resonance, why the sound is loud at this frequency.

[1 mark]



There is an increase in loudness when a stationary wave forms in the air in the tube.
This stationary wave corresponds to the first harmonic.

Figure 7 shows:

- the position **P** of a node at the surface of the water
- the position **Q** of an antinode.

Q is a constant distance x above the top of the tube.

0 7 . 2

Describe the motion of the air molecules at **P** and at **Q** due to the stationary wave.

[2 marks]

at **P** _____

at **Q** _____

The length L of the column of air is varied.

The frequency f of the first harmonic is recorded for each value of L .

0 7 . 3

When a stationary wave of the first harmonic is formed, the distance **PQ** is equal to $\frac{\lambda}{4}$, where λ is the wavelength of the stationary wave.

Show that f is given by $\frac{1}{f} = \frac{4L}{v} + \frac{4x}{v}$

where v is the speed of sound in air.

[1 mark]


Question 7 continues on the next page

Turn over ►



Figure 8 shows the variation of $\frac{1}{f}$ with L . The best-fit line has been drawn.

Figure 8



0 7 . 4 Determine v using **Figure 8**.

[3 marks]

$v =$ _____ m s^{-1}



0	7	.	5
---	---	---	---

Explain how you would use **Figure 8** to determine x .
Calculations are not required.

[2 marks]

9

Turn over for the next question

Turn over ►



0 8 . 1

Explain what is meant by modal dispersion in an optical fibre.

[2 marks]

The speed of light in the glass core of an optical fibre is $1.97 \times 10^8 \text{ m s}^{-1}$.

0 8 . 2

Calculate the refractive index of the glass.


[1 mark]

refractive index = _____

A pulse of light is passed along the optical fibre and experiences modal dispersion. The fibre is straight and has a length of 10.0 km.

The longest path of light along the fibre is at an angle θ_{max} to the central axis as shown in **Figure 9**.

The shortest path of light along the fibre is along the central axis.

Figure 9**not to scale**

There is a time delay of $13.7 \mu\text{s}$ between the arrival of the pulse when it follows the longest path and when it follows the shortest path.

0	8	.	3
---	---	---	---

Show that the length of the longest path is approximately 12.7 km.

[2 marks]

0	8	.	4
---	---	---	---

Calculate θ_{max} .

[1 mark]

$$\theta_{\text{max}} = \underline{\hspace{10em}}^{\circ}$$

Question 8 continues on the next page


Turn over ►



A laser sends a signal along the optical fibre. The signal is made up of a series of repeated pulses as shown in **Figure 10**.

Figure 10 shows the signal as it enters the fibre.

Figure 10



0 8 . 5

Calculate the frequency of repetition of the pulses.

[2 marks]

frequency = _____ Hz

0 8 . 6

Calculate the maximum duration of each pulse that is received at the end of this fibre.

[2 marks]

maximum duration = _____ s



0	8	.	7
---	---	---	---

Explain how pulse broadening limits the frequency at which pulses in a signal can be received through this optical fibre.

[1 mark]

11

Turn over for the next question

Turn over ►



0 9

In an experiment, a beam of electrons is accelerated from rest through a potential difference.

The beam is split into two coherent beams by a device that acts as a double slit. The two electron beams interfere to form an interference pattern as shown in **Figure 11**.


The path from the centre of the double slit to the n^{th} bright fringe makes an angle θ_n with the path to the central fringe.

The fringe separation is w .

The distance between the double slit and the interference pattern is D .

D is much larger than w .

Figure 11



0 9 . 1

Explain why θ_n in radians is approximately equal to $\frac{nw}{D}$.

[1 mark]



Figure 12 shows the interference pattern from this experiment.
The scale of the image is shown.

Figure 12



The slit separation of the double slit is 496 nm.

09.2

Show that the de Broglie wavelength of the electrons is approximately $1 \times 10^{-11} \text{ m}$.

In your answer:

- annotate **Figure 12** to show how you made your measurements
- use the relationship given in Question **09.1**.

[4 marks]

Question 9 continues on the next page

Turn over ►



0	9	.	3
---	---	---	---

Calculate the accelerating potential difference for the electrons.
Ignore relativistic effects.


[3 marks]

accelerating potential difference = _____ V

—
8

END OF SECTION A



Section BAnswer **all** questions in this section.**1 0****Figure 13** shows a diagram of an electric kettle.**Figure 13**

This kettle contains a heating filament that consists of coiled nichrome wire surrounded by an electrical insulator.

The heating filament is connected to the plug by a power cord that is made from copper wires surrounded by an electrical insulator.

When the kettle is switched on, energy is transferred from the filament to the water.

1 0 . 1

The resistance of the nichrome wire is $18\ \Omega$ at its operating temperature.
The voltage across the nichrome wire is $230\ \text{V}$.

Calculate the power transferred by the nichrome wire.

[1 mark]

power = _____ W

Question 10 continues on the next page**Turn over ►**

The resistance of the copper wire in the power cord must be much less than the resistance of the nichrome wire.

1 0 . 2

Explain why.

[3 marks]

1 0 . 3

The resistivity of nichrome is $1.12 \times 10^{-6} \Omega \text{ m}$ at its operating temperature.
The resistance of the nichrome wire is 18Ω at this temperature.
The radius of the nichrome wire is 0.16 mm.

Calculate the length of the nichrome wire.

[2 marks]

length = _____ m



1	0	.	4
---	---	---	---

When the kettle is switched on, the temperature of the nichrome wire increases rapidly for about 5 ms, until the nichrome wire reaches its operating temperature. The temperature then remains constant.

Explain how the power consumption of the kettle varies for the first 10 ms. Calculations are not required.

[3 marks]

9


Turn over for the next question

Turn over ►




1 1

Figure 14 shows an arrangement used to determine the wavelength of laser light. The light is incident normally on a diffraction grating. The diffraction pattern is viewed on a screen a distance D from the diffraction grating.

Figure 14

The distance between the two second-order maxima is x .

A student uses a metre ruler to measure x , as shown in **Figure 15**.

Figure 15

1 1 . 1

Determine x using **Figure 15**.
State an appropriate unit for your answer.

[1 mark]
 $x =$ _____

unit = _____



1 1 . 2

Calculate the percentage uncertainty in your value of x .**[1 mark]**

percentage uncertainty = _____

1 1 . 3

The student measures D as 200.0 cm.
The diffraction grating has 100 lines per mm.

Calculate, in nm, the wavelength λ of the laser light.**[3 marks]** $\lambda =$ _____ nm

1 1 . 4

The diffraction grating is now replaced with a grating that has a larger number of lines per mm. The uncertainty in the grating spacing is negligible.

Discuss how this will affect the percentage uncertainty in the measured value of λ .**[2 marks]**

END OF SECTION B

7
Turn over ►

Section C

Each of the questions in this section is followed by four responses, **A**, **B**, **C** and **D**.

For each question select the best response.

Only **one** answer per question is allowed.


For each question, completely fill in the circle alongside the appropriate answer.


CORRECT METHOD



WRONG METHODS



If you want to change your answer you must cross out your original answer as shown. 

If you wish to return to an answer previously crossed out, ring the answer you now wish to select as shown. 

You may do your working in the blank space around each question but this will not be marked.
Do **not** use additional pages for this working.

1 2 What is the unit for resistivity in SI fundamental (base) units?

[1 mark]

A $\text{kg m}^3 \text{s}^{-2} \text{A}^{-1}$

☐

B $\text{kg m}^3 \text{s}^{-3} \text{A}^{-1}$

☐

C $\text{kg m}^3 \text{s}^{-2} \text{A}^{-2}$


☐

D $\text{kg m}^3 \text{s}^{-3} \text{A}^{-2}$

☐


1 3

The circuit shown contains a battery of emf 12 V with negligible internal resistance.



What is the potential difference between **P** and **Q**?

[1 mark]

A 4 V

☐

B 6 V

☐

C 8 V

☐

D 12 V

☐


Turn over for the next question

Turn over ►



1 4

A potential difference is applied across a neon gas-discharge tube.



In the tube, neon ions (Ne^+) with a charge of $+e$ travel from the anode to the cathode. Free electrons move in the opposite direction.

In 3.0 minutes, 4.4×10^{20} Ne^+ ions arrive at the cathode and 4.4×10^{20} free electrons reach the anode.

What is the total current between the cathode and the anode?


[1 mark]

- A** 0 ☐
- B** 0.39 A ☐
- C** 0.78 A ☐
- D** 23 A ☐



1 5

In the circuit shown, the cell has negligible internal resistance.
T is a negative temperature coefficient thermistor.



The temperature of the thermistor changes, causing an increase in the current.

Which row shows the change in temperature and the effect this change has on the voltmeter reading?

[1 mark]

	Change in temperature	Effect on voltmeter reading	
A	increase	increase	<input type="radio"/>
B	increase	decrease	<input type="radio"/>
C	decrease	increase	<input type="radio"/>
D	decrease	decrease	<input type="radio"/>

Turn over for the next question

Turn over ►



Questions 16 and 17 are about the following experiment.

A student uses a stopwatch to investigate the time period of a pendulum.

There is an absolute uncertainty of 0.2 s at the beginning and the end of each timing.

1 6

The student measures ten oscillations of the pendulum. The stopwatch records the time for ten oscillations to be 15.07 s.

What is the best estimate for the percentage uncertainty of the time period of the pendulum?

[1 mark]

A 2.7%

☐

B 1.3%

☐

C 0.27%

☐

D 0.13%

☐

1 7

The student repeats the measurement three more times. His four measurements of the time for ten oscillations are:

15.07 s

15.35 s

15.02 s

14.92 s

What is the best estimate for the percentage uncertainty in this set of measurements?

[1 mark]

A 0.14%

☐

B 1.4%

☐

C 2.7%


☐

D 2.8%





☐

1 8

An oscillating mass–spring system has a time period T .
The graph shows the variation with time of the displacement of the oscillating mass.



Which graph shows the variation with time of potential energy E_p for the system?

[1 mark]**A****B****C****D**

Turn over for the next question


Turn over ►




1 9

A transverse wave is created by oscillating a string of beads.

Graph 1 shows the variation of the displacement of each bead with the distance measured from a fixed point along the wave.

Graph 1

Graph 2 shows the variation of the displacement of a particular bead with time.

Graph 2

Which row gives the wavelength and frequency of the wave?

[1 mark]

	Wavelength	Frequency	
A	K	H	<input type="radio"/>
B	K	$\frac{1}{H}$	<input type="radio"/>
C	$2K$	H	<input type="radio"/>
D	$2K$	$\frac{1}{H}$	<input type="radio"/>



2 0

Sound travels at a speed v in air and at a speed $4.5v$ in water.

A sound wave with a wavelength λ is produced in air. The wave passes from air into water.

Which row shows the wavelength and the nature of the wave in water?

[1 mark]

	Wavelength	Nature	
A	4.5λ	longitudinal	<input type="radio"/>
B	λ	longitudinal	<input type="radio"/>
C	4.5λ	transverse	<input type="radio"/>
D	λ	transverse	<input type="radio"/>

2 1

The frequency of the first harmonic of a violin string is 440 Hz.

The tension in the string is increased by 8%. The length and mass of the string do not change.

What is the new frequency of the first harmonic of the string?

[1 mark]

A 475 Hz ☐

B 457 Hz ☐

C 452 Hz ☐

D 422 Hz ☐

Turn over ►

2 2


Monochromatic light is incident normally on a diffraction grating.

The first-order maximum is observed at an angle of 18° to the zero-order maximum.

What is the largest angle at which a maximum can be found?

[1 mark]**A** 90° ☐**B** 72° ☐**C** 68° ☐**D** 54° ☐**2 3**

Light is incident on a Perspex–air boundary and is partially reflected and partially refracted. The incident and reflected rays are shown. The refracted ray is not shown.

**not to scale**

air

The reflected ray is deflected by a total angle of 36° from the incident ray. The refractive index of Perspex is 1.5

What is the angle of refraction in air for the refracted ray?

[1 mark]**A** 12° ☐**B** 23° ☐**C** 28° ☐**D** 62° ☐

2 4

A monochromatic light source **X** produces N_X photons per second, each with energy E_X .
 A monochromatic light source **Y** has the same power as **X** and emits light with a smaller frequency than **X**.
Y produces N_Y photons per second, each with energy E_Y .

Which row shows how N_X compares with N_Y and how E_X compares with E_Y ?

[1 mark]

A	$N_X < N_Y$	$E_X > E_Y$	<input type="radio"/>
B	$N_X < N_Y$	$E_X = E_Y$	<input type="radio"/>
C	$N_X = N_Y$	$E_X > E_Y$	<input type="radio"/>
D	$N_X = N_Y$	$E_X = E_Y$	<input type="radio"/>

Turn over for the next question


Turn over ►



2 5

Radiation of wavelength λ is incident on a metal surface. Photoelectrons are emitted with maximum kinetic energy E_k .

The graph shows how $\frac{1}{\lambda}$ varies with E_k .



What is the gradient of the graph?

[1 mark]

A h ☐

B hc ☐

C $\frac{1}{h}$ ☐

D $\frac{1}{hc}$ ☐

14

END OF QUESTIONS



There are no questions printed on this page

*Do not write
outside the
box*

**DO NOT WRITE ON THIS PAGE
ANSWER IN THE SPACES PROVIDED**



[illegible]

Do not write
outside the
box

[illegible]

